

# Homeoprophylaxis: Human Records, Studies and Trials

---

## In the beginning...

Homeopathy's ability to prevent as well as treat epidemic disease was first discovered in 1799 by Samuel Hahnemann, founder of Homeopathy, during an epidemic of scarlet fever in which he successfully used Belladonna to contain its spread in those he treated.

The prophylactic effects of Homeopathy were quickly picked up by other Homeopaths of the day and subsequently used with success in all of the great world epidemics. It is still employed in outbreaks and epidemics today - by people, communities and governments who have retained the knowledge and the will to use it. The following is a collation of some of those instances – historical and current.

## Viral Diseases

**2007:** Human herpesvirus 1, human adenovirus C serotype 5, influenza A virus, human respiratory syncytial virus, human parainfluenza virus 3, human rhinovirus B serotype 14, and human coxsackievirus serotype A9 cause, among other complaints, colds, flu, sore throat, runny nose, cold sores, bronchiolitis, pneumonia, hand foot and mouth disease, and conjunctivitis. Gripp-Heel, a proprietary combination remedy demonstrated significant in vitro reductions of infectivity by 20% to 40% when tested against these viruses. Gripp-Heel contains *Aconitum napellus*, *Byronia alba*, *Lachesis muta*, *Eupatorium perfoliatum* and *Phosphorus* in a range of decimal potencies.<sup>i</sup>

**2010:** Human rhinovirus B serotype 14, influenza A virus, H1N1 virus, herpes simplex virus 1, vesicular stomatitis virus, respiratory syncytial virus, parainfluenza type 3, and adenovirus cause, among other complaints, flu, colds, sore throat, swollen glands, oral vesicles, runny nose, cold sores, bronchiolitis, pneumonia, hand foot and mouth disease, and conjunctivitis. Proprietary products Engystol and Gripp-Heel displayed in vitro prophylactic effects when tested against these viruses. Gripp-Heel contains *Aconitum napellus*, *Byronia alba*, *Lachesis muta*, *Eupatorium perfoliatum* and *Phosphorus* in a range of decimal potencies. Engystol contains *Vincetoxicum hirundinaria* and *Sulphur* in a range of decimal potencies.<sup>ii</sup>

## Chikungunya

**2006:** In Kerala, India, a group of doctors distributed a homeopathic preventative called *Eupatorium perfoliatum* (Eup-per) for chikungunya. The results of the study showed that while over 73 %

contracted chikungunya in the unprotected group, only 17% of the protected group contracted the disease.<sup>iii</sup>

## Cholera

**1831:** Samuel Hahnemann prevented and treated cholera during the 1831 Asiatic cholera epidemic with the remedies Camphor, Cuprum metallicum and Veratrum album. Accounts can be read in Cure and Prevention of Asiatic Cholera.<sup>iv</sup>

**1849:** Dr Clemens von Boenninghausen treated and prevented “untold” numbers of cholera infections with the above remedies recommended by Hahnemann during the 1849 European epidemic. While a death rate of 54-90% occurred with conventional treatment, Boenninghausen’s patients had a mortality rate of only 5-16%.<sup>v</sup>

**1841-1854:** Dr Charge, after producing a statistical table showing the difference in cholera mortality between homeopathic and allopathic treatment at the Convent Refuge at Marseilles, stated that once homeopathy was introduced prophylactically, no further cases were reported.<sup>vi</sup>

## Dengue Fever

**1996:** The Central Council of Research in Homoeopathy reported that Dengueinum 30C was administered to at least 39,200 people in the Delhi area during an epidemic of dengue haemorrhagic fever. The follow-up of 23,520 people, 10 days later, revealed that only 5 people (0.125%) had developed mild symptoms, with the rest showing no signs or symptoms of the disease.<sup>vii</sup> (During epidemics of dengue, attack rates among the susceptible are often 40-50 %, but may reach 80-90 %, World Health Organisation)<sup>viii</sup>

**2001:** In São Paulo, Brazil in May 2001, a single dose of the homeopathic remedy Eupatorium perfoliatum 30C was given during a dengue outbreak to 40% of residents in the most highly affected neighbourhood. Thereafter, Dengue incidence decreased by 81.5%, a highly significant decrease when compared with those neighbourhoods that did not receive homeopathic prophylaxis ( $p < 0.0001$ ).<sup>ix</sup>

**2006:** A dengue outbreak was controlled by the Cuban Government through the preventative and treatment use of homeopathic remedies. A marked reduction in dengue haemorrhagic fever occurred easing the demand for intensive care beds.<sup>x</sup>

**2007:** Between April and September 2007, a homeopathic complex composed of Eupatorium perfoliatum, Phosphorus and Crotalus horridus 30C was given to 20,000 city residents in São Paulo, Brazil. Unfortunately this trial was aborted prematurely due to national political intervention, but the preliminary findings indicate that homeopathy was effective in the prevention and treatment of the dengue epidemic.<sup>xi</sup>

**2008:** Facing an outbreak of dengue in early 2007, the Secretary of Health of the county of Macaé, Rio de Janeiro, Brazil, carried out a “Homeopathy Campaign against Dengue”. 156,000 doses of the prophylactic remedy were freely distributed to asymptomatic patients and 129 doses of the ‘genus epidemicus’ to symptomatic patients within outpatient clinics. The remedy was a homeopathic complex containing Phosphorus 30C, Crotalus horridus 30C and Eupatorium perfoliatum 30C. The disease incidence in the first three months of 2008 fell 93% in comparison to the corresponding period in 2007. The rest of the untreated state experienced an increase of 128%.<sup>xii</sup>

**2008 – 2010:** An interview with the Secretary of Health responsible for the integrative homeopathy program of the previous paper reported the following: In 2007, facing a dengue epidemic, the city of Macaé, Rio de Janeiro, decided to add a homeopathic medicine to the contingency plan for dengue fever after which the disease incidence in Macaé in 2008 fell by 71% compared to 2007. In the north Fluminense region, which did not receive the prophylactic, there was an increase of 273%, and in the State of Rio de Janeiro, the increase was 315%. In 2011, there was a decrease of 89% of confirmed cases in the protected area in the first quarter compared to the same period in 2010. The prophylactic was freely distributed by the Health Department of Macaé in the years: 2007: 216,000 doses; 2008: 203,878 doses; 2009: 211,059 doses; 2010: 178 677 doses. In terms of lethality and notified dengue cases there was 0.2 in 2007, 0.0 in 2008, 0.1 in 2009 and 0.2 in 2010. In 2010 the agreed lethality target for severe forms in Macaé was 2.58; the municipality reached 0.68, a much lower rate compared to the state and other municipalities.<sup>xiii</sup>

**2010:** The Colombo Municipal Council (CMC) and homeopathic doctors used homeoprophylaxis to curb dengue in areas prone to dengue mosquito breeding in Colombo city and its suburbs.<sup>xiv</sup>

**2012:** In Mudurai, India, doctors from the government medical college and hospital dispensed the prophylactic Eupatorium perforatum to thousands during a dengue fever epidemic. First to receive the remedy were doctors, nurses, hospital staff and students followed by the residents of the city.<sup>xv</sup>

**2012:** The Government Homoeopathic Medical College and Hospital at Tirumangalam, India, distributed homeoprophylaxis for the prevention of dengue fever at two free camps during an outbreak.<sup>xvi</sup>

**2012:** The Medical and Health Department of Chittorr, India, distributed homeoprophylactics to 2.5 million people during a dengue fever epidemic following the earlier success of prophylactics distributed for the swine flu epidemic in Kurnool.<sup>xvii</sup>

**2012:** Researchers and doctors from KEM hospital used homeopathic Tabacum as the genus epidemicus for the treatment and prevention of dengue fever in Pune, India.<sup>xviii</sup>

**2012:** The Government Homoeopathic Medical College and Hospital at Tirumangalam, India, established more than 30 camps to distribute a homeoprophylactic for dengue fever control during regional outbreaks.<sup>xix</sup>

**2013:** AYUSH doctors in Guwahati state, India, used homeopathic remedies to treat and prevent dengue during a local outbreak. The former deputy director of health services and state programme

officer of AYUSH advised that the constitutional medicine and nosodes were among the best preventive medicines for the disease.<sup>xx</sup>

## Diphtheria

**1932:** Laboratory experiments published by Dr Chavanon showed that 45 children became Schick test negative (indicating the presence of antibodies to diphtheria) after being treated with Diphtherinum.<sup>xxi</sup>

**1941:** The test was repeated by Drs Patterson and Boyd with 23 out of 33 children becoming Schick test negative after being given Diphtherinum.<sup>xxii</sup>

**1947:** Dr Roux repeated the test and produced a similar result.<sup>xxiii</sup>

## Hepatitis

**1991:** Homeopathic Phosphorus has been recognised as a prophylactic for viral hepatitis for many years. Dr Eizayaga of Brazil noted both its curative and preventative effect. When treating individuals with hepatitis, he would also give their family members and school-mates the Phosphorus prophylactic. None contracted hepatitis in spite of being in close contact.<sup>xxiv</sup>

## Influenza and Respiratory Tract Infections

**1968:** Between 1968-70, a survey conducted in Indian factories and offices compared the results of allopathic (conventional) treatment and homeopathic treatment of influenza. The purpose of this survey was to determine the effectiveness of Influenzinum as a homeopathic preventative (prophylactic). Almost 20% of the patients treated by conventional medical physicians contracted the flu. Among the homeopathically treated patients, only 6.5% came down with the disease. Those who did become ill recovered more rapidly than their allopathically treated patients. The number of working days lost by the allopathically treated patients was nearly eight and a half times greater than those lost by homeopathic patients.<sup>xxv</sup>

**2007:** Human herpesvirus 1, human adenovirus C serotype 5, influenza A virus, human respiratory syncytial virus, human parainfluenza virus 3, human rhinovirus B serotype 14, and human coxsackievirus serotype A9 cause, among other complaints, colds, flu, sore throat, runny nose, cold sores, bronchiolitis, pneumonia, hand foot and mouth disease, and conjunctivitis. Gripp-Heel, a proprietary combination remedy demonstrated significant in vitro reductions of infectivity by 20% to 40% when tested against these viruses. Gripp-Heel contains Aconitum napellus, Byronia alba, Lachesis muta, Eupatorium perfoliatum and Phosphorus in a range of decimal potencies.<sup>xxvi</sup>

**2010:** Human rhinovirus B serotype 14, influenza A virus, H1N1 virus, herpes simplex virus 1, vesicular stomatitis virus, respiratory syncytial virus, parainfluenza type 3, and adenovirus cause,

among other complaints, flu, colds, sore throat, swollen glands, oral vesicles, runny nose, cold sores, bronchiolitis, pneumonia, hand foot and mouth disease, and conjunctivitis. Proprietary products Engystol and Gripp-Heel displayed in vitro prophylactic effects when tested against these viruses. Gripp-Heel contains Aconitum napellus, Byronia alba, Lachesis muta, Eupatorium perfoliatum and Phosphorus in a range of decimal potencies. Engystol contains Vincetoxicum hirundinaria and Sulphur in a range of decimal potencies.<sup>xxvii</sup>

**2011:** Nosodes of potential infecting agents, or placebo, were given for 30 days to 450 children in Brazil to test their effectiveness in preventing flu and acute respiratory infections. Over the next 12 months the incidence of diagnosed acute respiratory infection or flu was 3 times higher in the placebo group than those given the prophylactics. The researchers commented that the low cost of treatment and the absence of adverse effects made these nosodes a useful therapeutic option for the Brazilian Public Health Service.<sup>xxviii</sup>

## Japanese Encephalitis

**1999 - 2003:** Japanese encephalitis had caused significant morbidity and mortality in the state of Andhra Pradesh for several decades. In 1986, a high of 2038 cases with 638 deaths was recorded. Children were especially affected. A small decline in incidence occurred when homeopathic Belladonna was given as a prophylactic to some. When the Government Department of Indian Medicine and Homoeopathy began to systematically distribute prophylactics in 1999 via Public Health Centres, Government Homeopathic Dispensaries, homoeopathic pharmacies and homoeopathic doctors, notifications and mortality dropped to nil by 2003. The prophylactics were dispensed to approximately 20 million children under 15 years of age. The three prophylactics used were Belladonna, Calcarea carbonica, and Tuberculinum.<sup>xxix xxx</sup>

**2010:** Research from the School of Tropical Medicine, Kolkata, in collaboration with Central Council for Research in Homoeopathy (under the Department of AYUSH, Government of India) showed that Belladonna, in a range of homoeopathic potencies, was successful in controlling the virus in infected chick embryos. The placebo had no effect.<sup>xxxi</sup>

## Leptospirosis

**2007-8:** In late 2007, the Cuban government rapidly distributed a homeopathic nosode prepared from four leptospirosis strains to 2.3 million people at high risk of infection from an annual leptospirosis epidemic. The remaining population of 8.8 million was untreated. Within weeks, the treated provinces had an 84% decrease in disease incidence while the numbers of those infected in untreated provinces continued at expected historical levels. The intervention was “strongly associated with a drastic reduction of disease incidence resulting in complete control of the epidemic.” The protective effect continued into 2008 with an 84% reduction in leptospirosis cases for the treated areas though no further prophylactic had been given. Leptospirosis infections in untreated areas increased by 22%.<sup>xxxii</sup>

## Malaria

**2003:** A malaria trial conducted in Kenya between 2003-2005 used the Malaria co prophylactic with a group of 33 volunteers. Twenty-one of the volunteers had experienced 1 – 3 malaria episodes in the 18 months prior to the trial. During the trial, one person thought he may have developed malaria but this was not verified by blood test. After a full recovery in a matter of hours, malaria was considered unlikely. All other participants in the trial remained malaria-free.<sup>xxxiii</sup>

**2008:** At Kendu Bay, Kenya, where malaria is endemic, 34 clients participated in a homeoprophylaxis observational field study. In the 6 months prior to the study, all participants had experienced attacks of malaria or malaria-like symptoms at least once with 71% experiencing malaria or malaria-like symptoms at least once a month or every two weeks. Over the course of the study and use of the prophylactic only 12% (4 out of the 34 respondents) experienced an attack of malaria or malaria-like symptoms.<sup>xxxiv</sup>

**2011-2013:** Chhattisgarh, India is a known endemic area for malaria. Between 2004 -2010 it had an alarming increase in deaths from the disease. Determined to prevent a repeat or worsening in 2011 the Health Minister instructed that a known homeopathic treatment and prophylactic remedy, Chininum sulphuricum, be distributed to 9 areas within the affected state. The results of this initiative were "more than encouraging". Significant protection was achieved with data showing that higher the number of doses of Chininum sulphuricum the lower the number of malaria cases. Those in unprotected areas were 5 times more likely to contract the disease. On the strength of these results the program was expanded to 20 areas in 2012 and 40 areas in 2013.<sup>xxxv</sup>

**2013:** Researchers evaluated the efficacy of Cinchona officinalis 30C and Chelidonium majus 30C in combination therapy against lethal murine malaria in mice. The combination of remedies showed significant preventive activity with chemosuppression that was higher than the standard drug, pyrimethamine. It also showed a moderate curative activity with complete clearance of parasites in 50% of surviving mice. The researchers stated, "These findings point to the significant antiplasmodial efficacy of the combination of these homeopathic drugs against Plasmodium berghei." <sup>xxxvi</sup>

## Meningococcal Disease

**1974:** During a meningococcal epidemic in Brazil, 18,640 children were immunised homeopathically to protect against meningococcal infection, and 6,340 were not. The following results were obtained:

18,640 protected homeopathically – 4 cases of meningococcal infection.

6,340 not protected – 32 cases of meningococcal infection.

Based on the infection (attack) rate in the unprotected group, 94 cases of infection could have been expected in the homeopathically protected group. Instead, there were only four cases of

meningococcal infection, showing that the homeopathic option was 95% effective against the meningococcal disease.<sup>xxxvii</sup>

**1998:** The results of the first study (above) led to a larger government-funded study, 24 years later. It was conducted by two Professors of Medicine from the University Foundation in Blumenau, Brazil, and a Blumenau specialist physician and Health City Secretary. A total of 65,826 people between the ages of 0–20 were immunised homeopathically to protect against meningococcal disease while 23,532 were not. Over a 12 month period, the following results were obtained:

65,826 protected homeopathically – 4 cases of meningococcal infection.

23,532 not protected – 20 cases of meningococcal infection.

Based on the infection (attack) rate in the unprotected group, 58 cases of infection could have been expected in the homeopathically protected group. Instead, there were only four cases of meningococcal infection. Statistical analysis showed that homeopathic immunisation offered 95% protection in the first six months and 91% protection over the year against meningococcal disease.<sup>xxxviii</sup>

## Pertussis

See 'Whooping Cough'.

## Poliomyelitis

**1800s:** Dr Grimmer of Chicago prophylactically treated 5,000 young children with *Lathyrus sativus*. None developed polio.<sup>xxxix</sup>

**1850:** During an epidemic of poliomyelitis, Dr Taylor Smith of Johannesburg, South Africa protected 82 people with homoeopathic *Lathyrus sativus*. Of these people, 12 came into direct contact with disease. None were infected.<sup>xl</sup>

**1956:** In a study between 1956-58, Dr Eisfelder immunized over 6,000 children with *Lathyrus sativus*. No cases of polio were reported in the group and no side-effects were observed.<sup>xli</sup>

**1957:** A severe poliomyelitis epidemic occurred in Buenos Aires. The majority of homoeopathic doctors prescribed *Lathyrus sativus* as a preventative. Drug stores distributed thousands of doses to the public. None of who used the prophylactic registered a case of contagion.<sup>xlii</sup>

**1975:** During another poliomyelitis epidemic in Buenos Aires, 40,000 were given the homeopathic prophylactic *Lathyrus sativus*. None developed poliomyelitis.<sup>xliii</sup>

## Scarlet Fever

**1799:** Homeoprophylaxis was born during an epidemic of Scarlet Fever in Germany. In 1799 Samuel Hahnemann, founder of Homeopathy, discovered during an epidemic that homeopathic Belladonna prevented Scarlet Fever in that particular epidemic if given to people before exposure.<sup>xliv</sup>

**1799:** Following Hahnemann's example (above), another eleven medical doctors prescribed Belladonna during the same epidemic. They reported that of 1,646 children exposed to scarlet fever after being given Belladonna, only 123 (7.4%) developed symptoms of infection. In contrast, the infection rate in those who did not receive the prophylactic was as high as 90%.<sup>xlv</sup>

**1838:** The Prussian Government ordered the use of Belladonna during all scarlet fever epidemics after a report from their protomedicus, Hufeland, showed it to be an effective prophylactic.<sup>xlvi</sup>

## Smallpox

**1800s:** Clemens von Boenninghausen (1785-1864) used Thuja in both the treatment and prevention of smallpox during an epidemic. When given to uninfected family members of households with members already sick with the disease, none went on to contract it.<sup>xlvii</sup>

**1902:** Dr Eaton reported that during a smallpox epidemic in Iowa, 2806 patients were treated prophylactically with Variolinum. Of the 547 patients definitely exposed, only 14 developed the disease. The protection rate from these numbers was 97%.<sup>xlviii</sup>

## Whooping Cough

**1967:** Dr Dorothy Shepherd provides several accounts of whooping cough prophylaxis by Pertussin. The following is one example: "Personally I have always used for years and years in whooping cough its own nosode, Pertussin, which Dr John Clarke recommended so strongly in his monograph. Pertussin has given me 100% protection even though it raged in a particular district; all the children on Pertussin, either on the 12th or 30th potency, given daily, escaped it. I used it for years in the nurseries, in my private practice, in the medical clinic, and in hundreds of cases in various epidemics we went through, and truly it is a great remedy."<sup>xlix</sup>

**1987:** A pilot study on whooping cough prophylaxis conducted by Dr J. English reported a reduction in whooping cough by a factor of approximately one in three when compared to the incidence of whooping cough in a similar but unprotected group.<sup>i</sup>

**1987:** A small clinical trial on prophylaxis with Pertussin by Dr A. Fox over a five-year period received 61 parent responses in which 5% of children definitely contracted whooping cough and 82% remained whooping cough free (the other 13% developed undiagnosed coughs). The prophylaxis was well tolerated with 67% of parents rating its effectiveness as "very good".<sup>ii</sup>

**2004:** Dr Golden compared the pertussis attack rate between unimmunised children and those given the whooping cough prophylactic during his 15-year study. From this he was able to determine efficacy. The unimmunised had an attack rate of 85% while the homeopathically protected group had an attack rate of 11.7%, producing a figure of efficacy of 86.2%.<sup>lii</sup>

**2012:** In an Irish trial involving 112 children 1 -2 months of age, those given homeoprophylactics for childhood diseases, including whooping cough, were less likely to experience severe symptoms of the diseases if contracted than the vaccinated comparison group.<sup>liii</sup>

---

<sup>i</sup> Glatthaar-Saalmuller B (2007). In vitro evaluation of the antiviral effects of the homeopathic preparation Gripp-Heel on selected respiratory viruses. Canadian Journal of Physiology and Pharmacology; 85(11): 1084-1090.

<sup>ii</sup> Roeska, K and Seilheimer, B. Journal of Immune Based Therapies and Vaccines 2010, 8:6  
<http://www.jibtherapies.com/content/8/1/6>

<sup>iii</sup> Dr R Rejikumar, Dr R S Dinesh et al. A Study on the Prophylactic Efficacy of Homoeopathic Preventive Medicine Against Chikungunya Fever. <http://www.similima.com/pdf/efficacy-chiunguna-kerala.pdf>

<sup>iv</sup> Hahnemann S. 1801. Cure and Prevention of Asiatic Cholera, p.755, The Lesser Writings of Samuel Hahnemann, republished by B. Jain Publishers, 1993, New Dehli.

<sup>v</sup> Von Boenninghausen, C. Baron.1984. Bönninghausens Kleine medizinische Schriften [Lesser Medical Writings] (ed. Klaus H. Gypser), Heidelberg, 1984.

<sup>vi</sup> Bradford, T. L., (Compiled by) 1900. The Logic of Figures or Comparative Results of Homeopathic and Other Treatments. p.130. Philadelphia: Boericke and Tafel. Reprinted by Nabu Public Domain Reprints. 2010. (Original Bradford source: North American Journal of Homeopathy, volume 4, p.298.)

<sup>vii</sup> Central Council of Research in Homoeopathy. CCRH News 1996-1997.

<sup>viii</sup> WHO Fact Sheet: Dengue and Dengue Haemorrhagic Fever <http://www.who.int/inf-fs/en/fact117.html>

<sup>ix</sup> Marino R (2008). Homeopathy and collective health: The case of dengue epidemics. International Journal of High Dilution Research; 7(25): 179-185.

<sup>x</sup> Secretaria Municipal de Saúde e Higiene da Prefeitura de São José do Rio Preto. Informe Técnico – 21/03/2007. 2007.

<sup>xi</sup> Marino R (2008). Homeopathy and collective health: The case of dengue epidemics. International Journal of High Dilution Research; 7(25): 179-185.

(<http://www.feg.unesp.br/~ojs/index.php/ijhdr/article/viewFile/312/373>)

<sup>xii</sup> de Souza Nunes LA (2008). Contribution of homeopathy to the control of an outbreak of dengue in Macaé, Rio de Janeiro. International Journal of High Dilution Research; 7(25): 186-192.

<sup>xiii</sup> Dengue and Homeopathy: a successful experience from Macae  
<http://www.ecomedicina.com.br/site/conteudo/entrevista22.asp>

<sup>xiv</sup> Sri Lanka Daily News: Homeopathy to Treat Dengue <http://archives.dailynews.lk/2010/08/04/news52.asp>

<sup>xv</sup> The Hindu: Keep Dengue at Bay with Homeopathy  
<http://www.thehindu.com/news/cities/Madurai/article3531685.ece>

<sup>xvi</sup> The Hindu: Homeopathy Camp for Dengue <http://www.thehindu.com/todays-paper/tp-national/tp-tamilnadu/article3553076.ece>

<sup>xvii</sup> The Hindu: Homeopathy to fight dengue fever <http://www.thehindu.com/todays-paper/tp-national/tp-andhrapradesh/homeopathy-to-fight-dengue-fever/article3767663.ece>

<sup>xviii</sup> Homeopathy is new challenger for Dengue  
<http://www.punemirror.in/article/2/20121110201211100842483927ea199fd/Homeopathy-is-new-challenger-for-dengue.html>

<sup>xix</sup> The Hindu: Dengue fever- homoeopathy camps evoke good response <http://www.thehindu.com/todays-paper/tp-national/tp-tamilnadu/dengue-fever-homoeopathy-camps-evoke-good-response/article4155839.ece>

- 
- <sup>xx</sup> The Times of India: Dengue menace - Homeopathic remedy explored  
[http://articles.timesofindia.indiatimes.com/2013-09-12/guwahati/42007941\\_1\\_dengue-menace-dengue-fever-dengue-cases](http://articles.timesofindia.indiatimes.com/2013-09-12/guwahati/42007941_1_dengue-menace-dengue-fever-dengue-cases)
- <sup>xxi</sup> Chavanon, P. 1952. *La Dipterie*, 4th Ed, St Denis, Niort: Imprimerie.
- <sup>xxii</sup> Patterson, J. and Boyd, W.E. Potency Action: A Preliminary Study of the Alternation of the Schick Test by a Homeopathic Potency. *British Homeopathic Journal*. 1941; 31: pp.301-309.
- <sup>xxiii</sup> Eizayaga. F. *Tratamiento Homeopatico de las Enfermedades Agudas y Su Prevencion*. Homeopatia. 1985; 51(324): pp. 352-362.
- <sup>xxiv</sup> *Treatise on Homeopathic Medicine* by Francisco Eizayaga, MD, published by Ediciones Maracel, Buenos Aires, Brazil, 1991.
- <sup>xxv</sup> Krishnamurty, Report on the use of Influenzinum during the outbreak of epidemic in India in 1968. *Hahnemannian Gleanings* 1970;37:225-6.
- <sup>xxvi</sup> Glatthaar-Saalmuller B (2007). In vitro evaluation of the antiviral effects of the homeopathic preparation Gripp-Heel on selected respiratory viruses. *Canadian Journal of Physiology and Pharmacology*; 85(11): 1084-1090.
- <sup>xxvii</sup> Roeska, K and Seilheimer, B. *Journal of Immune Based Therapies and Vaccines* 2010, 8:6  
<http://www.jibtherapies.com/content/8/1/6>
- <sup>xxviii</sup> Lyrio C, Siqueira CM, Veiga VF, Homsani F, Marques AL, Biolchini J, Dantas F, de Matos HJ, Passos SRL, Couceiro JN, Holandino C (2011). The use of homeopathy to prevent symptoms of human flu and acute respiratory infections: A double-blind, randomized, placebo-controlled clinical trial with 600 children from Brazilian public health service. *International Journal of High Dilution Research*; 10(36): 174-176.  
<http://www.feg.unesp.br/~ojs/index.php/ijhdr/article/view/499/513>
- <sup>xxix</sup> The Hindu: Japanese Encephalitis on the decline in State  
<http://www.hindu.com/2003/04/02/stories/2003040204970400.htm>
- <sup>xxx</sup> Golden, I. 2010. *Vaccination and Homeoprophylaxis: A Review of the Risks and Alternatives*. 7th Edition. Isaac Golden Publications.
- <sup>xxxi</sup> Bandyopadhyay B, Das S, Sengupta M, Saha C, Das KC, Sarkar D, Nayak C (2010). Decreased intensity of Japanese encephalitis virus infection in chick chorioallantoic membrane under influence of ultradiluted belladonna extract. *American Journal of Infectious Diseases*; 6(2): 24-28.
- <sup>xxxii</sup> Bracho G, Varela E, Fernández R, et al. Large-scale application of highly-diluted bacteria for Leptospirosis epidemic control. *Homeopathy* 2010; 99: 156-166.
- <sup>xxxiii</sup> Partington, T. Silent and Deadly: Prophylaxis and Treatment of Malaria. *Homeopathy in Practice*, 2006, pp.14-19.
- <sup>xxxiv</sup> The Use of Homeopathic Prophylaxis and Treatment For Malaria in Endemic Areas Of Kenya (Part 2)  
<http://hpathy.com/homeopathy-papers/the-use-of-homeopathic-prophylaxis-and-treatment-for-malaria-in-endemic-areas-of-kenya/>
- <sup>xxxv</sup> Efficacy of Chininum Sulph 200C in prevention of malaria - An Operational Research Study. State Health Resource Centre, Chhattisgarh.  
[http://www.shsrc.org/Downloads/Chininum%20-Final%20-\(17June13\).pdf](http://www.shsrc.org/Downloads/Chininum%20-Final%20-(17June13).pdf)
- <sup>xxxvi</sup> Rajan A, Bagai U. Antimalarial potential of China 30 and Chelidonium 30 in combination therapy against lethal rodent malaria parasite: Plasmodium berghei. *Journal of Complementary & Integrative Medicine*.  
<http://www.ncbi.nlm.nih.gov/pubmed/23652641>
- <sup>xxxvii</sup> Castro, D. & Nogueira, G. G. (1975). Use of the nosode Meningococcinum as a preventative against meningitis. *Journal of the American Institute of Homeopathy*, 1975 Dec 68 (4), 211-219.
- <sup>xxxviii</sup> Mronisnski C, Adriano E & Mattos G. (1998/99) Meningococcinum: Its protective effect against Meningococcal disease, *Homeopathic Links*, Vol 14 Winter 2001, 230-234.
- <sup>xxxix</sup> Currim, A.M. Ed. 1996. *The Collected Works of Arthur Grimmer, M.D.* Norwalk and Greifenberg: Hahnemann International Institute for Homeopathic Documentation.
- <sup>xl</sup> Taylor-Smith, A. "Poliomyelitis and prophylaxis. *British Homeopathic Journal*, 1950 Apr;40(2):65-77
- <sup>xli</sup> Eisfelder, H. "Poliomyelitis Immunization: A Final Report." *Journal of the American Institute of Homeopathy*. V. 54, Nov-Dec 1961, pp. 166-167.
-

- 
- <sup>xlii</sup> Treatise on Homeopathic Medicine by Francisco Eizayaga, MD, published by Ediciones Maracel, Buenos Aires, Brazil, 1991.
- <sup>xliii</sup> Treatise on Homeopathic Medicine by Francisco Eizayaga, MD, published by Ediciones Maracel, Buenos Aires, Brazil, 1991.
- <sup>xliv</sup> Hahnemann S. 1801. Cure and Prevention of Scarlet Fever, p. 369-384, The Lesser Writings of Samuel Hahnemann, republished by B. Jain Publishers, 1993, New Dehli.
- <sup>xlv</sup> Dudgeon R. E. 1853. Lectures on the Theory and Practice of Homeopathy, p.541,2. Republished by B. Jain Publishers, 2002, New Dehli.
- <sup>xlvi</sup> Hufeland. 1828. Prophylactic Power of Belladonna in Scarlet Fever, Hufeland's Journal.
- <sup>xlvii</sup> Von Boenninghausen, C. Baron. Vide Concerning the Curative Effects of Thuja in Smallpox. Lesser Writings.
- <sup>xlviii</sup> Eaton, C.W. Variolinum. Transactions of the American Institute of Homeopathy, 1907. P. 547-567.
- <sup>xlix</sup> Shepherd, D., (1967). Homeopathy in epidemic diseases (First ed.). Essex, England: The C. W. Daniel Company Limited. p.18.
- <sup>i</sup> English, J.M. "Pertussin 30—Preventive for whooping cough? A pilot study," The British Homeopathic Journal, April 1987, Vol. 76, p. 61--65.
- <sup>ii</sup> Fox, A. D. (1987). Whooping cough prophylaxis with pertussin 30. British Homoeopathic Journal, 76(2), 69-70.
- <sup>iii</sup> Golden, I. 2010. Vaccination and Homeoprophylaxis: A Review of the Risks and Alternatives. 7th Edition. p.194. Isaac Golden Publications.
- <sup>liii</sup> Claudia, D. R., Ciara, O. R., Sarah, O. B., & Edward, S. (2012). Homeopathic prophylaxis: suggestion for Vaccination Homeopathic Protocol. European Journal of Integrative Medicine, 4, 36.